


For parents of children with ADHD



## Control the symptoms. Lose the label. Find his true potential.

**ADDERALL XR®** may make a difference in his life, by reducing ADHD symptoms to a level similar to that of children without ADHD—in one week.

- 87% of parents preferred **ADDERALL XR** over the medication their child was taking before
- Each convenient, once-daily dose begins working quickly and keeps working into homework time and family time
- The active ingredient in **ADDERALL XR** has a 65-year history of medical use

Ask your doctor about  
**ADDERALL XR.** It can make  
a difference for your child.

[adderallxr.com](http://adderallxr.com) 1-800 000 0000



**5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg CAPSULES**  
(Mixed Salts of a Single-Entity Amphetamine Product)  
Dextroamphetamine Sulfate Dextroamphetamine Saccharate  
Amphetamine Aspartate Monohydrate Amphetamine Sulfate

**IMPORTANT SAFETY INFORMATION:** Adderall XR was generally well tolerated in clinical studies. The most common side effects in studies included: *children*—decreased appetite, difficulty falling asleep, stomachache, and emotional lability; *adolescents*—loss of appetite, difficulty falling asleep, stomachache, and weight loss; *adults*—dry mouth, loss of appetite, difficulty falling asleep, headache, and weight loss.

Adderall XR may not be right for everyone. Patients should speak with their doctor if they have a history of high blood pressure or any heart conditions, glaucoma, thyroid problems, emotional instability, mental illness, or a known allergy to this type of medication. Abuse of amphetamines may lead to dependence. Misuse of amphetamines may cause sudden death and serious cardiovascular adverse events. These events have also been reported rarely with amphetamine use.

If you are currently taking or have recently taken a type of antidepressant called a MAO inhibitor or have a pre-existing structural heart abnormality, you should not take Adderall XR. There is a potential for worsening of motion or verbal tics and Tourette's syndrome. A patient should report any new psychological symptoms to his or her physician.

Please see brief summary of prescribing information on adjacent page. For more information, consult your physician.

**Shire** ...your ADHD Support Company™

©2005 Shire US Inc., Wayne, Pennsylvania 19087 AD00X XX00X

For parents of children with ADHD

# Control the symptoms. Lose the label. Find her true potential.

**ADDERALL XR®** may make a difference in his life, by reducing ADHD symptoms to a level similar to that of children without ADHD—in one week.

- **87% of parents preferred ADDERALL XR over the medication their child was taking before**
- Each convenient, once-daily dose begins working quickly and keeps working into homework time and family time
- The active ingredient in **ADDERALL XR** has a 65-year history of medical use

ONE DOSE DAILY

**ADDERALL XR®** (II)

**5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg CAPSULES**  
(Mixed Salts of a Single-Entity Amphetamine Product)  
Dextroamphetamine Sulfate, Dextroamphetamine Saccharate,  
Amphetamine Aspartate Monohydrate, Amphetamine Sulfate

**Ask your doctor about ADDERALL XR.  
It can make a difference for  
your child.**

**adderallexr.com 1-800 000 0000**

**IMPORTANT SAFETY INFORMATION:** Adderall XR was generally well tolerated in clinical studies. The most common side effects in studies included: *children*—decreased appetite, difficulty falling asleep, stomachache, and emotional lability; *adolescents*—loss of appetite, difficulty falling asleep, stomachache, and weight loss; *adults*—dry mouth, loss of appetite, difficulty falling asleep, headache, and weight loss.

Adderall XR may not be right for everyone. Patients should speak with their doctor if they have a history of high blood pressure or any heart conditions, glaucoma, thyroid problems, emotional instability, mental illness, or a known allergy to this type of medication. Abuse of amphetamines may lead to dependence. Misuse of amphetamines may cause sudden death and serious cardiovascular adverse events. These events have also been reported rarely with amphetamine use.

If you are currently taking or have recently taken a type of antidepressant called a MAO inhibitor or have a pre-existing structural heart abnormality, you should not take Adderall XR. There is a potential for worsening of motion or verbal tics and Tourette's syndrome. A patient should report any new psychological symptoms to his or her physician.

Please see brief summary of prescribing information on adjacent page. For more information, consult your physician.

**Shire** ...your ADHD Support Company™